Review Article

Medicinal Plants for Improving Cognitive Function: A Review

Behnam Sadighi^{1#}, Ghasem Khazen^{2#}, Somaiyeh Taheri-Targhi³, Zahra Yousefi^{4*}, Mostafa Araj-Khodaei^{5*}

¹Medicinal Plants Research Center, Maragheh University of Medical Sciences, Maragheh, Iran

²Depertment of Persian Medicine, Faculty of Traditional Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

³Aging Research Institute, Tabriz University of Medical Sciences, Tabriz, Iran

⁴Research Center of Psychiatry and Behavioral Sciences, Aging Research Institute, Tabriz University of Medical Sciences, Tabriz Iran

Tabriz, Iran

⁵Research Center for Integrative Medicine in Aging, Aging Research Institute, Tabriz University of Medical Sciences, Tabriz, Iran

Article History: Received: October 12, 2024 Accepted: November 2, 2024 ePublished: November 26, 2024

***Corresponding Authors:** Zahra Yousefi,

Email: zahra69_y@yahoo.com and Mostafa Araj-Khodaei, Email: mostafaa33@gmail.com

*Contributed equally as the first author.

Abstract

Background: Cognition is critical for functional independence as people age. Neurodegenerative dementias are more likely to affect older adults due to the aging process. The inadequate response of conventional therapeutic approaches, besides numerous side effects, can cause patients to explore complementary and alternative medical options. This study aimed to introduce the most common medicinal herbs used in Persian medicine for the prevention and treatment of cognitive impairments.

Methods: In order to identify medicinal herbs for the treatment and prevention of cognitive diseases, manuscripts were searched by keywords such as "Nesyan", "Fesad-al-Zekr", and "Zekr". Afterwards, the scientific names and pharmacological properties of selected plants were also searched in scientific databases such as PubMed, Scopus, and Web of Science.

Results: The results of the review indicate that herbal remedies could be effective in treating cognitive impairment and dementia. The neuroprotective, antioxidant, and acetylcholinesterase inhibitory activities are the basis for how these remedies work.

Conclusion: The neuropharmacological effects of medicinal plants on numerous pathways make them promising candidates for future drugs or adjunct therapies in combination with established pharmaceuticals; however, further clinical research is required to establish the safety and effectiveness of these treatments.

Keywords: Medicinal plants, Cognition, Review

Please cite this article as follows: Sadighi B, Khazen G, Taheri-Targhi S, Yousefi Z, Araj-Khodaei M. Medicinal plants for improving cognitive function: a review. Int J Drug Res Clin. 2024; 2: e22. doi: 10.34172/ijdrc.2024.e22

Introduction

Owing to the improvement of health care in the last decades, the number of the elderly has increased in most societies.1 Among other common diseases in individuals aged 65 years and above, cognitive impairment is one of the most prevalent dysfunctions. It is a neurodegenerative disorder that affects different abilities such as memory, attention, and executive functioning.² language, Neurodegenerative and vascular diseases, and dysthymia or dysphoria can result in memory impairment and are usually a sign of dementia. The age-adjusted prevalence of dementia in the Middle East and North Africa region was 777.6 per 100000 people in 2019, representing an increase of 3% compared to 1990.3 According to various surveys, the prevalence of the disease varies in different populations. The prevalence of Alzheimer's disease and mild cognitive impairment in the elderly aged 65 years and

above was reported to be 11.3% and 22.7%, respectively.4 It is predicted that the number of individuals with memory impairment will double by 2040 in developing countries.5 The disease not only reduces life satisfaction in patients but also imposes a huge burden on society and health care. Considering the rapid growth in the prevalence of the disease and problems that are imposed on both patients and society, the development of new methods of treatment and prevention seems necessary.⁶ There is currently no cure for dementia but certain medications can help manage symptoms of the disease and slow down the worsening of symptoms. Even though the drugs used for the treatment of cognitive impairment are generally well tolerated, with common side effects being nausea, diarrhea, and vertigo, there is a requirement for more alternative drugs that have fewer side effects and focus on the underlying neurobiological mechanisms.7



© 2024 The Author(s). This is an open access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Numerous herbal remedies have been used in the prevention and treatment of cognitive impairment in traditional medicine.⁸ Recently, different studies have evaluated the efficiency of herbal remedies in treating chronic neurodegenerative disorders. The current study introduces the most common medicinal herbs for the prevention and treatment of memory impairment in Persian Medicine. Furthermore, this study supports the evidence of pharmacological properties of recommended herbs.

Materials and Methods

Ghanoon fel-teb, Gharabadin Kabir, and Exir-e-Azam are the main Persian traditional medicine references. These books have been taught in European medical schools for centuries. Manuscripts authored by eminent scientists across centuries were chosen. The selected books were searched using keywords such as "Nesyan", "Fesad-al-Zekr", and "Zekr" to find the related medicinal herbs for the treatment and prevention of diseases related to cognition. After gathering the data from old manuscripts, the scientific names and pharmacological properties of selected plants were also searched in scientific databases such as PubMed, Scopus, and Web of Science.

Studies that explored the use of plants for the prevention and treatment of cognitive impairment (animal, human, and in vitro) and were published in authentic journals were selected. Research articles from journals with uncertain validity or without a clear research strategy were excluded from the study (Table 1).

Lavandula angustifolia

Lavandula angustifolia, native to the Mediterranean area, belongs to the Lamiaceae family. Different parts of the plant such as leaves and flowers either in the form of oil or other extracts are used as therapeutic agents in Traditional Medicine. Besides the sedative, anticonvulsant, analgesic, and local anesthetic effects, the aqueous extract of the plant has beneficial impacts on memory (Lavender). Caffeic acid and luteolin-7-glycosid are the two constituents of the aqueous extract of lavender that are potentially effective in reducing oxidative stress and inflammatory response; therefore, this plant can be useful in the treatment of neurodegenerative diseases.54 Based on the results of a study conducted in China, lavender was efficient as an adjunct treatment in relieving agitated behaviors of Chinese patients who suffered from dementia. Therefore, among the patients that are particularly vulnerable to side effects of psychotropic medications, alternative solutions such as aromatherapy by lavender could be a good choice for both patients and clinicians.55

Ferula assa-foetida

Ferula assa-foetida is an herbaceous perennial plant of the Apiaceae family that grows widely in the central and southern mountains of Iran. Asafoetida is an oleo-gumresin obtained from the exudates of the roots of the *F*. *assa-foetida*. Because of its sulfurous odor and bitter taste, it is used as a flavoring spice in Iranian cuisine. This plant has been reported to be beneficial for the brain and nerves due to inhibition of AChE activity that results in the improvement of memory and learning. The secondary metabolites in plant extracts are able to increase the amount of acetylcholine in the brain so it can alleviate the symptoms of neurological dysfunction in individuals with Alzheimer's disease or dementia.⁵⁶

Terminalia chebula Retz

Terminalia chebula Retz, known as black myrobalan, is a medicinal plant whose different parts such as fruit, bark, and root are used in the treatment of various ailments. The plant belongs to the *Combretaceae* family. Based on the findings of several surveys, the fruits of the plant are useful in the improvement of memory loss and cognitive deficits. Additionally, *T. chebula Retz* contains large amounts of active compounds which have beneficial effects on Alzheimer's disease.⁵⁷

Allium sativum L.

Allium sativum L., known as garlic, is one of the most common herbs used both as flavor and medicine since ancient times. According to different studies, it has anti-amyloidogenic, anti-oxidant, and acetylcholine activities, which can prevent common age-related diseases.⁵⁸

Melissa officinalis

Melissa officinalis, which is commonly referred to as "lemon balm" due to its lemon-like flavor and fragrance, is a member of the *Lamiaceae* family. The plant has been used as medicine traditionally in many parts of the world. Among its various functions, the anti-depressant effect is of considerable interest.⁵⁹ The terpenes, esters and aldehydes present in the lemon balm produce antioxidant and sedative effects. The essential oil of the plant has been traditionally used for relieving distress and anxiety for centuries.⁶⁰

Thymus vulgaris

Thymus vulgaris is a flowering plant that is native to southern Europe from the western Mediterranean to southern Italy and belongs to the Lamiaceae family. The plant not only is used as medicine but also as a culinary ingredient and aroma. Due to its antioxidant and acetylcholinesterase inhibitory effects, it is useful in the treatment of dementia.⁶¹

As mentioned before, the best therapeutic approach for the treatment of dementia is the inhibition of AChE. Based on the results of several surveys, *Thymus vulgaris* had a neuroprotective effect on scopolamine-induced memory impairment in zebrafish. It can be concluded that the plant has an important role in improving the cholinergic nervous system and antioxidative stress. Therefore, it can be a good alternative approach for the treatment of cognitive dysfunction.²⁰

Table 1	Pharmacological	Properties of Most	Important Medicinal	Plants Used for the	e Treatment of	Dementia in TPM

Scientific Name of the plant	Traditional Name	Part Used	Pharmacological Effect	Dose	Animal/ in vitro/ RCT	Reference
Lavandula angustifolia	Ustukhuddus	Arial part	Neuroprotective, anti- Alzheimer, antioxidant, antispasmodic, and acetylcholinesterase inhibitory effects	1-2 g	Animal/in vitro/RCT	9,10
Ferula assa- foetida	Anjedan	Seed	Acetylcholinesterase inhibitory effect and antioxidant properties	200-500 g	Animal/in vitro	11,12
Terminalia chebula Retz	Halileh kaboli	Fruit	Neuroprotective, anti- Alzheimer, antioxidant, and acetylcholinesterase inhibitory effects	500-1500 g	Animal/in vitro	13,14
Allium sativum L.	Som	Bulb	Anti-amyloidogenics, antioxidant, and acetylcholinesterase inhibitory effects	300-1000 g	Animal/in vitro/RCT	15,16
Melissa officinalis L.	Badranjbuyeh	Aerial part	Neuroprotective, anti- Alzheimer, antioxidant, and acetylcholinesterase inhibitory effects	1-2 g	Animal/in vitro/RCT	17,18
Thymus vulgaris	Hasha	Leaf	Neuroprotective, antioxidant, and acetylcholinesterase inhibitory effects	3-4 g	Animal/ in vitro/ RCT	12,19,20
Peganum harmala. L.	Harmal	Seed	Cerebroprotective effect, acetylcholinesterase inhibitory effect, suppressing lipid peroxidation, augmenting endogenous antioxidant enzymes, and reducing acetylcholinesterase activity in the brain ²¹	2-4 g	Animal/in vitro	11,12
Pimpinella anisum L.	Anisun	Fruit	Neuroprotective and antioxidant effects and reducing both AChE and oxidative stress	1-1.5 g	Animal/in vitro	22,23
Rosa damascena	Gol-e Sorkh	Flower	Antioxidant effect, induce neurogenesis, synaptic plasticity, improve memory improve significantly the levels of acetylcholinesterase enzymes and the rate of stress indicators, and improving the level of consciousness and brain function	1.5 g	In vitro	24,25
Hyssopus officinalis L.	Zofa	Arial part	No significant acetylcholinesterase inhibitory effect, antioxidant effect	450-900 g	Animal/in vitro	26,27
<i>Pistacia lentiscus</i> L.	Mastaki	Gum	Neuroprotective, anti- Alzheimer, antioxidant, and acetylcholinesterase inhibitory effects	200 g	Animal/in vitro	28,29
Piper nigrum L.	Felfel	Fruit	Neuroprotective, anti- Alzheimer, and antioxidant effects, reducing AChE activity	4.5 g	Animal/in vitro	30-32
Anacyclus pyrethrum L.	Aghergherha	Root	Neuroprotective, anti- Alzheimer, antioxidant, and acetylcholinesterase inhibitory effects	0.5-1 g	Animal/in vitro	33,34
Acorus calamus L.	Vaj	Root	Neuroprotective, anti- Alzheimer, antioxidant, and acetylcholinesterase inhibitory effects	4.5 g	Animal/in vitro	35,36
Crocus sativus L.	Zaferan	Stigma	Neuroprotective, anti- Alzheimer, and antioxidant effects	20-400 mg	Animal/in vitro/RCT	37-39
Teucrium polium L	Jade	Leaf	Anticonvulsant, antioxidant, anti-Alzheimer, and anti-inflammatory activities	250 mg	Animal/in vitro	12,40
Cinnamomum verum	Darchin	Stem bark	Neuroprotective, anti- Alzheimer, and antioxidant activities	2.5 g	Animal/in vitro	29,41
Zingiber officinale Rosc	Zanjabil	Rhizome	Anti-inflammatory, anti-apoptotic, neuroprotective, anti-Alzheimer, and antioxidant activities	70-140 mg	Animal/in vitro	42-44
Boswellia carterii	Kondor	Oleo- gum-resin	Neuroprotective, anti- Alzheimer, antioxidant, and acetylcholinesterase inhibitory effects	300-400 mg	Animal/in vitro	29,45,46
Piper longum	Darfilfl	Fruit	Neuroprotective, anti- Alzheimer, and antioxidant activities	0.5-1 g	Animal/in vitro/RCT	47,48
Cyperus rotundus.	So'd	Tuber	Neuroprotective effects such as neurogenesis and neuronal regeneration that improve learning and memory performance, anti-Alzheimer activity, and antioxidant activity	3-6 g	In vitro	49,50
Phyllanthus emblica L.	Amele	Fruit	Neuroprotective, anti- Alzheimer, antioxidant, activities, decrease in the level of AChE activity	250-500 g	Animal/in vitro	45,51
Asarum europaeum	Osaron	Rhizome	Neuroprotective, anti- Alzheimer, antioxidant, and acetylcholinesterase inhibitory effects	4.5 g	In vitro	45,52
Terminalia bellirica	Balile	Fruit	Neuroprotective, anti- Alzheimer, and antioxidant activities	500-1500 mg	Animal/in vitro	13
Vitis vinifera	Zabib	Fruit	Neuroprotective, anti- Alzheimer, and antioxidant activities	4-18 mL	Animal/in vitro/RCT	53

Peganum harmala L.

Peganum harmala L. is a perennial herb whose seeds and aerial parts have been used as medicine in China for the treatment of various diseases such as forgetfulness for centuries. In a study, Liu et al demonstrated that different parts of the plant are used for treating cognitive dysfunctions such as Alzheimer's disease. 62

The two β -carboline alkaloids, harmine and harmaline,

are responsible for AChE inhibition. In summary, the plant extract may have a significant role in the prevention of dementia. The inhibition of oxidative stress increases the activity of antioxidant enzymes which results in improvement of enzyme activities in the brain.¹¹

Pimpinella anisum L.

Pimpinella anisum L. is a medicinal plant that belongs to warm regions like India, Iran, and Egypt. Different studies have shown that the plant has therapeutic effects on several ailments such as neurologic disorders.

According to the reports of different studies, there are several mechanisms like scavenging free radicals and neuroprotection that are caused by inhibition of acetylcholinesterase in brain cells and result in antiamnesic action.^{63,64}

Rosa damascena

Rosa damascena is a bushy shrub with colorful flowers used in different industries such as perfume, food, and medicine. The origin of the plant is the Middle East but it is cultivated all over the world. The medicinal function of the plant is attributed to the presence of phenolic compounds that have anti-oxidant activity which improves memory and brain function.⁶⁵

Hyssopus officinalis L.

Hyssopus officinalis L. belongs to the *Lamiaceae* family and is distributed from the East Mediterranean to Central Asia. *Hyssopus officinalis* has been used as a medicinal remedy for a long time in different nations. One of the activities of *H. officinalis* proved by surveys is its antioxidant function.⁶⁶

Pistacia lentiscus L.

Pistacia lentiscus L. is a member of the *Anacardiaceae* family. The most commonly used part of the plant is resin which is known as Mastic resin and exudes from the bark. The plant has antiatherogenic, antimicrobial, antioxidant, hepatoprotective, antiarthritic, wound healing, and anticancer properties.⁶⁷

Piper nigrum L.

Piper nigrum L., known as pepper, is the most commonly consumed spice in the world. Almost all parts of the plant such as flower, seed, leaf, and fruit are used for managing different ailments. According to different in vitro and in vivo studies, the fruit of the plant ameliorates memory impairment by attenuating oxidative stress in rat hippocampus.⁶⁸

Anacyclus pyrethrum L.

Anacyclus pyrethrum L., a plant native to Asian countries such as India, belongs to the Asteraceae family. The roots of the plant are used for therapeutic purposes. Based on the reports of various surveys, the plant has memory-enhancing activity and is efficient in the treatment of impaired cognitive functions.⁶⁹ The results of a study demonstrated that the ethanolic extract of *A*. *pyrethrum* has memory-enhancing effect by increasing cholinesterase level in the brain. Therefore, it improves cognitive processes.⁷⁰

Acorus calamus L.

Acorus calamus is a medicinal herb from the *Acoraceae* family. The rhizome of the plant when dried and powdered has a spicy flavor. Rhizome and leaves of the plant have also been used as medicine traditionally.

Crocus sativus L.

Crocus sativus L., known as Saffron, is widely cultivated in Iran and other countries such as India and Greece. Saffron has been traditionally used for the treatment of different diseases like depression.⁷¹ The pistil of the plant is useful in the treatment of mental illnesses and improvement of memory.⁷² *Acorus calamus* reduces oxidative damage induced by scopolamine, thereby restoring the brain's antioxidant state. Therefore, it may be an efficient therapeutic option for the treatment and prevention of neurodegenerative disorders like dementia and Alzheimer's disease.^{73,74}

Teucrium polium L.

Teucrium polium L., a member of the *Lamiaceae* family, can be found abundantly in Iran and grows wildly. Among different properties of the plant such as hyperglycemic and insulinotropic activities, some reports have demonstrated the antioxidant effect of the herb as well.⁷⁵ Results show that *T. polium* is rich in essential minerals like calcium, sodium, magnesium, and so on. Moreover, the phenolic compounds of the plant have antioxidant, antihyperglycaemic, anti-Alzheimer, and anti-inflammatory properties. Therefore, it can be a good candidate for alleviating different health-debilitating problems.⁴⁰

Cinnamomum verum

Cinnamon (*Cinnamomum verum*), grown in Sri Lanka and Southern India, is a traditional herb with various health benefits that has been used for the treatment of different ailments.⁷⁶ Research has demonstrated that cinnamon has been traditionally utilized for agerelated brain disorders. Potent BChE inhibitory activity of cinnamon can be considered in the management of advanced Alzheimer's disease (AD). The polyphenolic compounds in the aqueous extract of cinnamon are responsible for its predominant antioxidant activity.⁴¹

Zingiber officinale Rosc.

Ginger, which is botanically known as *Zingiber officinale* Roscoe, is a medicinal herb from the Zingiberaceae family. The phenolic compounds of ginger such as gingerols and shogaols are the primary agents responsible for the health benefits of ginger.⁷⁷ The antioxidant and anti-acetylcholinesterase activities of ginger have been studied for the effectiveness of ginger in preventing and treating degenerative diseases, including Alzheimer's disease.⁴² Studies have shown that ginger can hinder the progression of dementia in all stages, neurodegeneration and neuroinflammation, and simultaneously maintain the survival of neurons.⁴²

Boswellia carterii

Boswellia carterii, known as Kondor, or Frankincense gum resin, is used in traditional medicine in India and Iran.⁷⁸ It is a valuable substance used in traditional medicine for its effect on the improvement of memory and learning. Avicenna, a Persian physician, proposed that *B. carterii* could improve memory function and prevent amnesia in the elderly.⁷⁹ The anti-oxidative, anti-inflammatory, and anti-diabetic properties, as well as its inhibitory effects on AChE and anti-A β activities, with its effects on the modulation of Tau deposition and hyperphosphorylation in the brain, make it a potent multi-targeting agent.⁷⁸

Piper longum

Although *Piper longum* is mostly used as a spice in cooking, several studies have proved the efficacy of the plant in the treatment of many diseases. Among various properties of the herb, its therapeutic potential for Alzheimer's disease is noticeable. A survey conducted by Joen et al demonstrated that the mixture of two ingredients, piperlonguminine and dihydrpiperlongominine, can control the expression of amyloid precursor protein which is helpful in managing the patients with Alzheimer's disease.⁸⁰

Cyperus rotundus

Cyperus rotundus belongs to the Cyperaceae family and due to its adaptability to wide range of climate and soil conditions, it can be cultivated in many parts of the world. Many studies revealed the antioxidant activity of the herb which is beneficial for managing loss of memory.⁸¹ The bioactive compounds of the plant such as phenolics and flavonoids are responsible for its antioxidant and antiapoptotic activities.⁸²

Phyllanthus emblica L.

Phyllanthus emblica is widely distributed in tropical areas such as Malaysia, India, Indonesia, and China. The delicious fruit of the plant is a great source of vitamin C and superoxide and has been traditionally used as medicine. Based on reports provided by different surveys, the phenolic compounds of the plant act as antioxidants and protect cells from the oxidation caused by free radicals. Free radicals are among the main causes of some neurodegenerative diseases.⁸³

Asarum europaeum

Asarum europaeum L. is a member of the Aristolochiaceae family. The plant has been used as a medicinal remedy for centuries in many countries. In Persian traditional

medicine, the rhizome of the plant has been used for the treatment of memory impairment. The results of the study conducted by Limon et al demonstrated that the neuroprotectivity of the components against A β in the hippocampus resulted in a significant improvement in spatial memory in rats.⁵²

Terminalia bellirica

Terminalia bellirica belongs to the *Combretaceae* family. This large tree is widely distributed in many parts of the world, especially in the Indian subcontinent and South East Asia. Oxidative stress is associated with many disorders in humans. Antioxidants inhibit or reduce the reaction of free radicals that protects cells against damage.⁸⁴ Several studies have reported the antioxidant activity of the plant, which is attributed to the presence of large amounts of phenolic and flavonoid components in the plant.⁸⁵

Vitis vinifera

Vitis vinifera is a well-known grape species of the Vitaceae family. There are seedless and non-seedless varieties in red, black, and white colors. It can be found in Western Asia and Southern Europe. All parts of the plant such as roots, stems, leaves, fruits, seeds, cane, and pomace contain different phytochemical substances. The ethanol extract of grape seeds has the antioxidant activity.⁸⁶ According to the results of a previous study, polyphenols derived from grapes have anti-oxidative, anti-inflammatory, and anti-amyloidogenic activities. Therefore, it can be considered a potential candidate for counteracting the multifactorial nature of Alzheimer's disease.⁸⁷

Discussion

As mentioned before, cognitive impairment/dementia is one of the most serious dysfunctions with a high prevalence among the elderly. Dementia is a neurodegenerative disorder that is chronic and progressive in nature. Several functions such as thinking, judgement, comprehension, learning, and calculation can be impaired by the disease.

The categorization of different types of cognitive impairment or dementia dates back to AD 1025 when Avicenna described the disease in his medical masterpiece "Canon of Medicine". Ibn Sina, known as Avicenna, was born on August 23, 980, in a small city of Iran, Afshaneh. His great talent for learning made him a genius in many subjects such as mathematics, astronomy, geometry, and theology when he was just a child. By the age of 17, he found medicine appealing and started the practice of medicine.⁸⁸ Qanun fi Al-Tibb was his encyclopedia of medicine and it overshadowed all previous medicinal books when appeared.⁸⁹ It became an important reference book in many universities of the West and was translated into many languages such as Latin, German, French, and so on.

Generally, cognitive impairment is defined as "Nesyan" in the third volume of Canon of Medicine. According to
 Table 2. The Most Important Activities of the Introduced Remedies

Mechanism	Name of the plant
Neuroprotective activity	Lavandula angustifolia, Terminalia chebula Retz, Melissa officinalis L., Melissa officinalis L., Thymus vulgaris, Pimpinella anisum L., Pistacia lentiscus L., Piper nigrum L., Anacyclus pyrethrum L., Acorus calamus L., Crocus sativus L, Cinnamomum verum, Zingiber officinale Rosc, Boswellia carterii, Piper longum, Piper longum, Cyperus rotundus, Phyllanthus emblica L., Asarum europaeum, Terminalia bellirica, Vitis vinifera
Antioxidant activity	Lavandula angustifolia, Ferula assa-foetida, Terminalia chebula Retz, Allium sativum L., Melissa officinalis L., Thymus vulgaris, Peganum harmala. L., Rosa damascene, Pimpinella anisum L., Hyssopus officinalis L., Pistacia lentiscus L., Piper nigrum L., Anacyclus, pyrethrum L., Acorus calamus L., Crocus sativus L., Teucrium polium L., Cinnamomum verum, Zingiber officinale Rosc, Boswellia carterii, Piper longum, Cyperus rotundus, Phyllanthus emblica L., Asarum europaeum, Terminalia bellirica, Vitis vinifera
Acetylcholinesterase inhibitory effect	Lavandula angustifolia, Ferula assa-foetida, Terminalia chebula Retz, Allium sativum L., Melissa officinalis L., Thymus vulgaris, Peganum harmala. L., Rosa damascene, Pistacia lentiscus L., Anacyclus pyrethrum L., Acorus calamus L., Asarum europaeum

him, different brain damages are caused by changes in brain temperament.

There are three main approaches to the treatment and prevention of ailments in traditional medicine:

- Lifestyle modification
- Manipulation
- Consumption of herbal remedies

Lifestyle Modification

According to Avicenna, the main goal of medicine is to maintain the well-being of individuals or return it when an illness occurs. Avicenna was the first scholar who explained the important factors associated with lifestyle.⁹⁰ The six principles of lifestyle modification include:

- Climate
- Mental movement and repose
- Sleep and wakefulness
- Foods and drinks
- Excretion of unnecessary substances and retention of necessary ones
- Body movement and repose

Manipulation

Manual therapy is the next step towards the treatment of diseases in traditional medicine. It has different types and indications. Massage is one form of manual intervention used by many scholars of traditional medicine.⁹¹

Herbal Remedies

Medicinal plants are precious sources of natural components that have been proven to play important roles as natural pharmaceutical agents. The use of plants as medicine dates back to the beginning of life on the earth by mankind. Various pharmacological studies and clinical trials have reported the efficacy of herbal remedies in the prevention and treatment of neurodegenerative disorders like cognitive impairment and dementia. It can be concluded that leaves are the most common part of the plant used for the treatment of cognitive impairment. However, it should be mentioned that other parts of the plants like rhizome, fruit, and seed are efficient in the prevention and treatment of the disorder as well. Based on the results of various surveys, several efficient mechanisms for memory improvement have been reported.

The most notable properties of the introduced remedies have been categorized in Table 2.

Conclusion

Based on the significance of chronic diseases like cognitive impairment and dementia and considering the high costs of treatment and the burden on both the healthcare system and society, it seems that further investigations are necessary as these remedies are natural and easy to use.

Ethics statement

Not applicable.

Disclosure of funding source

None.

Conflict of interests declaration

The authors declare no conflict of interests.

Data availability statement

Not applicable.

Author contributions

Conceptualization: Behnam Sadighi, Mostafa Araj-Khodaei. Investigation: Zahra Yousefi, Mostafa Araj-Khodaei, Ghasem Khazen. Methodology: Mostafa Araj-Khodaei, Zahra Yousefi. Supervision: Mostafa Araj-Khodaei. Writing-original draft: Zahra Yousefi, Somaiyeh Taheri-Targhi, Ghasem Khazen.

Writing-review & editing: Behnam Sadighi, Mostafa Araj-Khodaei.

Consent for publication

Not applicable.

References

- Norton S, Matthews FE, Barnes DE, Yaffe K, Brayne C. Potential for primary prevention of Alzheimer's disease: an analysis of population-based data. Lancet Neurol. 2014;13(8):788-94. doi: 10.1016/s1474-4422(14)70136-x.
- Bai W, Chen P, Cai H, Zhang Q, Su Z, Cheung T, et al. Worldwide prevalence of mild cognitive impairment among community dwellers aged 50 years and older: a meta-analysis and systematic review of epidemiology studies. Age Ageing. 2022;51(8):afac173. doi: 10.1093/ageing/afac173.
- Zarei M, Asheghi M, Zarei M. Artificial intelligence in dementia diagnosis: past, present, and future. Int J Aging. 2023;1:e14. doi: 10.34172/ija.2023.e14.
- Fratiglioni L, De Ronchi D, Agüero-Torres H. Worldwide prevalence and incidence of dementia. Drugs Aging. 1999;15(5):365-75. doi: 10.2165/00002512-199915050-00004.
- Prince M, Wimo A, Guerchet M, Ali GC, Wu YT, Prina M. World Alzheimer Report 2015. The Global Impact of Dementia: An Analysis of Prevalence, Incidence, Cost and Trends. London: Alzheimer's Disease International; 2015.

- Safiri S, Noori M, Nejadghaderi SA, Mousavi SE, Sullman MJ, Araj-Khodaei M, et al. The burden of Alzheimer's disease and other types of dementia in the Middle East and North Africa region, 1990-2019. Age Ageing. 2023;52(3):afad042. doi: 10.1093/ageing/afad042.
- Benninghoff J, Perneczky R. Anti-dementia medications and anti-Alzheimer's disease drugs: side effects, contraindications, and interactions. In: Riederer P, Laux G, Nagatsu T, Le W, Riederer C, eds. NeuroPsychopharmacotherapy. Cham: Springer; 2022. p. 2779-88. doi: 10.1007/978-3-030-62059-2_195.
- Kim HG, Oh MS. Herbal medicines for the prevention and treatment of Alzheimer's disease. Curr Pharm Des. 2012;18(1):57-75.
- Hancianu M, Cioanca O, Mihasan M, Hritcu L. Neuroprotective effects of inhaled lavender oil on scopolamine-induced dementia via anti-oxidative activities in rats. Phytomedicine. 2013;20(5):446-52. doi: 10.1016/j.phymed.2012.12.005.
- Spiridon I, Colceru S, Anghel N, Teaca CA, Bodirlau R, Armatu A. Antioxidant capacity and total phenolic contents of oregano (*Origanum vulgare*), lavender (*Lavandula angustifolia*) and lemon balm (*Melissa officinalis*) from Romania. Nat Prod Res. 2011;25(17):1657-61. doi: 10.1080/14786419.2010.521502.
- Adhami HR, Farsam H, Krenn L. Screening of medicinal plants from Iranian traditional medicine for acetylcholinesterase inhibition. Phytother Res. 2011;25(8):1148-52. doi: 10.1002/ ptr.3409.
- Shirbeigi L, Dalfardi B, Abolhassanzadeh Z, Nejatbakhsh F. Dementia etiologies and remedies in traditional Persian medicine; a review of medicinal plants and phytochemistry. Curr Drug Metab. 2018;19(5):414-23. doi: 10.2174/1389200 218666170810170124.
- Afshari AR, Sadeghnia HR, Mollazadeh H. A review on potential mechanisms of *Terminalia chebula* in Alzheimer's disease. Adv Pharmacol Sci. 2016;2016:8964849. doi: 10.1155/2016/8964849.
- 14. Chang CL, Lin CS. Phytochemical composition, antioxidant activity, and neuroprotective effect of *Terminalia chebula* Retzius extracts. Evid Based Complement Alternat Med. 2012;2012:125247. doi: 10.1155/2012/125247.
- Tasnim S, Haque PS, Bari MS, Hossain MM, Islam SM, Shahriar M, et al. *Allium sativum* L. improves visual memory and attention in healthy human volunteers. Evid Based Complement Alternat Med. 2015;2015:103416. doi: 10.1155/2015/103416.
- 16. Chauhan NB. Anti-amyloidogenic effect of *Allium sativum* in Alzheimer's transgenic model Tg2576. J Herb Pharmacother. 2003;3(1):95-107.
- 17. Koksal E, Bursal E, Dikici E, Tozoglu F, Gulcin I. Antioxidant activity of *Melissa officinalis* leaves. J Med Plants Res. 2011;5(2):217-22.
- Ferreira A, Proença C, Serralheiro ML, Araújo ME. The in vitro screening for acetylcholinesterase inhibition and antioxidant activity of medicinal plants from Portugal. J Ethnopharmacol. 2006;108(1):31-7. doi: 10.1016/j.jep.2006.04.010.
- Rabiei Z, Mokhtari S, Asgharzade S, Gholami M, Rahnama S, Rafieian-Kopaei M. Inhibitory effect of *Thymus vulgaris* extract on memory impairment induced by scopolamine in rat. Asian Pac J Trop Biomed. 2015;5(10):845-51. doi: 10.1016/j. apjtb.2015.07.006.
- 20. Capatina L, Todirascu-Ciornea E, Napoli EM, Ruberto G, Hritcu L, Dumitru G. *Thymus vulgaris* essential oil protects zebrafish against cognitive dysfunction by regulating cholinergic and antioxidants systems. Antioxidants (Basel). 2020;9(11):1083. doi: 10.3390/antiox9111083.
- 21. Osman NN, Alanbari KH, Al-Shreef HA. Evaluation of the possible antioxidant effects of *Peganum harmala* and *Ginkgo biloba* in ameliorating Alzheimer's disease in rat model.

Int J Pharm Sci Res. 2018;9(8):3189-98. doi: 10.13040/ ijpsr.0975-8232.9(8).3189-98.

- Mushtaq A, Anwar R, Ahmad M. Methanolic extract of *Pimpinella anisum* L. prevents dementia by reducing oxidative stress in neuronal pathways of hypermnesic mice. Pak J Zool. 2020;52(5):1779-86. doi: 10.17582/journal. pjz/20181002061042.
- 23. Tepe B, Akpulat HA, Sokmen M, Daferera D, Yumrutas O, Aydin E, et al. Screening of the antioxidative and antimicrobial properties of the essential oils of *Pimpinella anisetum* and *Pimpinella flabellifolia* from Turkey. Food Chem. 2006;97(4):719-24. doi: 10.1016/j.foodchem.2005.05.045.
- 24. Beigom Hejaziyan L, Hosseini SM, Taravati A, Asadi M, Bakhshi M, Moshaei Nezhad P, et al. Effect of *Rosa damascena* extract on rat model Alzheimer's disease: a histopathological, behavioral, enzyme activities, and oxidative stress study. Evid Based Complement Alternat Med. 2023;2023:4926151. doi: 10.1155/2023/4926151.
- Moure A, Franco D, Sineiro J, Domínguez H, Núñez MJ, Lema JM. Antioxidant activity of extracts from *Gevuina avellana* and *Rosa rubiginosa* defatted seeds. Food Res Int. 2001;34(2-3):103-9. doi: 10.1016/s0963-9969(00)00136-8.
- Alinezhad H, Azimi R, Zare M, Ebrahimzadeh MA, Eslami S, Nabavi SF, et al. Antioxidant and antihemolytic activities of ethanolic extract of flowers, leaves, and stems of *Hyssopus officinalis* L. Var. angustifolius. Int J Food Prop. 2013;16(5):1169-78. doi: 10.1080/10942912.2011.578319.
- Fathiazad F, Hamedeyazdan S. A review on *Hyssopus* officinalis L.: composition and biological activities. Afr J Pharm Pharmacol. 2011;5(17):1959-66. doi: 10.5897/ajpp11.527.
- Quartu M, Serra MP, Boi M, Pillolla G, Melis T, Poddighe L, et al. Effect of acute administration of *Pistacia lentiscus* L. essential oil on rat cerebral cortex following transient bilateral common carotid artery occlusion. Lipids Health Dis. 2012;11:8. doi: 10.1186/1476-511x-11-8.
- 29. Yousefsani BS, Barreto GE, Sahebkar A. Beneficial medicinal plants for memory and cognitive functions based on traditional Persian medicine. Adv Exp Med Biol. 2021;1308:283-90. doi: 10.1007/978-3-030-64872-5_20.
- Farzaei MH, Shahpiri Z, Mehri MR, Bahramsoltani R, Rezaei M, Raeesdana A, et al. Medicinal plants in neurodegenerative diseases: perspective of traditional Persian medicine. Curr Drug Metab. 2018;19(5):429-42. doi: 10.2174/1389200219 666180305150256.
- Hritcu L, Noumedem JA, Cioanca O, Hancianu M, Kuete V, Mihasan M. Methanolic extract of *Piper nigrum* fruits improves memory impairment by decreasing brain oxidative stress in amyloid beta(1-42) rat model of Alzheimer's disease. Cell Mol Neurobiol. 2014;34(3):437-49. doi: 10.1007/s10571-014-0028-y.
- Yang W, Chen YH, Liu H, Qu HD. Neuroprotective effects of piperine on the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridineinduced Parkinson's disease mouse model. Int J Mol Med. 2015;36(5):1369-76. doi: 10.3892/ijmm.2015.2356.
- Selles C, El Amine Dib M, Allali H, Tabti B. Evaluation of antimicrobial and antioxidant activities of solvent extracts of *Anacyclus pyrethrum* L., from Algeria. Mediterr J Chem. 2012;2(2):408-15. doi: 10.13171/mjc.2.2.2012.01.11.22.
- Sujith K, Ronald DC, Suba V. Inhibitory effect of *Anacyclus* pyrethrum extract on acetylcholinesterase enzyme by invitro methods. Pharmacogn J. 2012;4(33):31-4. doi: 10.5530/ pj.2012.33.5.
- 35. Ahmed F, Chandra JN, Urooj A, Rangappa KS. In vitro antioxidant and anticholinesterase activity of *Acorus calamus* and *Nardostachys jatamansi* rhizomes. J Pharm Res. 2009;2(5):830-83.
- 36. Muthuraman A, Singh N. Attenuating effect of Acorus calamus extract in chronic constriction injury induced neuropathic

pain in rats: an evidence of anti-oxidative, anti-inflammatory, neuroprotective and calcium inhibitory effects. BMC Complement Altern Med. 2011;11:24. doi: 10.1186/1472-6882-11-24.

- Saleem S, Ahmad M, Ahmad AS, Yousuf S, Ansari MA, Khan MB, et al. Effect of saffron (*Crocus sativus*) on neurobehavioral and neurochemical changes in cerebral ischemia in rats. J Med Food. 2006;9(2):246-53. doi: 10.1089/jmf.2006.9.246.
- Akhondzadeh S, Shafiee Sabet M, Harirchian MH, Togha M, Cheraghmakani H, Razeghi S, et al. A 22-week, multicenter, randomized, double-blind controlled trial of *Crocus sativus* in the treatment of mild-to-moderate Alzheimer's disease. Psychopharmacology (Berl). 2010;207(4):637-43. doi: 10.1007/s00213-009-1706-1.
- 39. Karimi E, Oskoueian E, Hendra R, Jaafar HZ. Evaluation of *Crocus sativus* L. stigma phenolic and flavonoid compounds and its antioxidant activity. Molecules. 2010;15(9):6244-56. doi: 10.3390/molecules15096244.
- Benchikha N, Messaoudi M, Larkem I, Ouakouak H, Rebiai A, Boubekeur S, et al. Evaluation of possible antioxidant, antihyperglycaemic, anti-Alzheimer and anti-inflammatory effects of *Teucrium polium* aerial parts (Lamiaceae). Life (Basel). 2022;12(10):1579. doi: 10.3390/life12101579.
- Ramshini H, Ebrahim-Habibi A, Aryanejad S, Rad A. Effect of *Cinnamomum verum* extract on the amyloid formation of hen egg-white lysozyme and study of its possible role in Alzheimer's disease. Basic Clin Neurosci. 2015;6(1):29-37.
- 42. Jagtap SR, Pol SL, Bhosale SS, Kadam VJ. Memory enhancing activity of ginger (*Zingiber officinale*), its treatments in dementia and Alzheimer's disease. Int J Res Appl Sci Biotechnol. 2022;9(3):73-84. doi: 10.31033/ijrasb.9.3.14.
- 43. Mathew M, Subramanian S. In vitro evaluation of anti-Alzheimer effects of dry ginger (*Zingiber officinale* Roscoe) extract. Indian J Exp Biol. 2014;52(6):606-12.
- 44. Waggas AM. Neuroprotective evaluation of extract of ginger (*Zingiber officinale*) root in monosodium glutamate-induced toxicity in different brain areas male albino rats. Pak J Biol Sci. 2009;12(3):201-12. doi: 10.3923/pjbs.2009.201.212.
- Jamshidi AH, Eghbalian F, Mahroozade S, Mohammadi Kenari H, Ghobadi A, Yousefsani BS. Recommended natural products in Alzheimer's disease based on traditional Persian medicine. J Med Plants. 2020;19(75):17-29. doi: 10.29252/ jmp.19.75.17. [Persian].
- Hosseinzadeh H, Ramezani M, Akhtar Y, Ziaei T. Effects Boswellia carterii gum resin fractions on intact memory and hyoscine-induced learning impairments in rats performing the Morris water maze task. J Med Plants. 2010;9(34):95-101. [Persian].
- Reddy NJ, Nagoor Vali D, Rani M, Rani SS. Evaluation of antioxidant, antibacterial and cytotoxic effects of green synthesized silver nanoparticles by *Piper longum* fruit. Mater Sci Eng C Mater Biol Appl. 2014;34:115-22. doi: 10.1016/j. msec.2013.08.039.
- Bi Y, Qu PC, Wang QS, Zheng L, Liu HL, Luo R, et al. Neuroprotective effects of alkaloids from *Piper longum* in a MPTP-inducedmousemodel of Parkinson's disease. Pharm Biol. 2015;53(10):1516-24. doi: 10.3109/13880209.2014.991835.
- 49. Shakerin Z, Esfandiari E, Razavi S, Alaei H, Ghanadian M, Dashti G. Effects of *Cyperus rotundus* extract on spatial memory impairment and neuronal differentiation in rat model of Alzheimer's disease. Adv Biomed Res. 2020;9:17. doi: 10.4103/abr.abr_173_19.
- Mohammadi M, Kazemitabar K, Asili J, Kamali H. Study of the antioxidant and antibacterial activity in methanolic, dichloromethan and hexane extracts of aerial parts of *Cyperus longos*. J North Khorasan Univ Med Sci. 2014;6(1):161-7. doi: 10.29252/jnkums.6.1.161. [Persian].
- 51. Uddin MS, Mamun AA, Hossain MS, Akter F, Iqbal MA,

Asaduzzaman M. Exploring the effect of *Phyllanthus emblica* L. on cognitive performance, brain antioxidant markers and acetylcholinesterase activity in rats: promising natural gift for the mitigation of Alzheimer's disease. Ann Neurosci. 2016;23(4):218-29. doi: 10.1159/000449482.

- 52. Saeedi M, Vahedi-Mazdabadi Y, Rastegari A, Soleimani M, Eftekhari M, Akbarzadeh T, et al. Evaluation of *Asarum europaeum* L. rhizome for the biological activities related to Alzheimer's disease. Res J Pharmacogn. 2020;7(3):25-33. doi: 10.22127/rjp.2020.223143.1565.
- 53. Jayaprakasha GK, Singh RP, Sakariah KK. Antioxidant activity of grape seed (*Vitis vinifera*) extracts on peroxidation models in vitro. Food Chem. 2001;73(3):285-90. doi: 10.1016/s0308-8146(00)00298-3.
- Soheili M, Salami M. Lavandula angustifolia biological characteristics: an in vitro study. J Cell Physiol. 2019;234(9):16424-30. doi: 10.1002/jcp.28311.
- 55. Lin PW, Chan WC, Ng BF, Lam LC. Efficacy of aromatherapy (*Lavandula angustifolia*) as an intervention for agitated behaviours in Chinese older persons with dementia: a cross-over randomized trial. Int J Geriatr Psychiatry. 2007;22(5):405-10. doi: 10.1002/gps.1688.
- Iranshahy M, Iranshahi M. Traditional uses, phytochemistry and pharmacology of asafoetida (*Ferula assa-foetida* oleogum-resin)-a review. J Ethnopharmacol. 2011;134(1):1-10. doi: 10.1016/j.jep.2010.11.067.
- 57. Kumar R, Arora R, Agarwal A, Gupta YK. Protective effect of *Terminalia chebula* against seizures, seizure-induced cognitive impairment and oxidative stress in experimental models of seizures in rats. J Ethnopharmacol. 2018;215:124-31. doi: 10.1016/j.jep.2017.12.008.
- Capasso A. Antioxidant action and therapeutic efficacy of *Allium sativum* L. Molecules. 2013;18(1):690-700. doi: 10.3390/molecules18010690.
- 59. Araj-Khodaei M, Noorbala AA, Yarani R, Emadi F, Emaratkar E, Faghihzadeh S, et al. A double-blind, randomized pilot study for comparison of *Melissa officinalis* L. and *Lavandula angustifolia* Mill. with fluoxetine for the treatment of depression. BMC Complement Med Ther. 2020;20(1):207. doi: 10.1186/s12906-020-03003-5.
- Moradkhani H, Sargsyan E, Bibak H, Naseri B, Sadat-Hosseini M, Fayazi-Barjin A, et al. *Melissa officinalis* L., a valuable medicine plant. J Med Plants Res. 2010;4(25):2753-9.
- 61. Prasanth Reddy V, Ravi Vital K, Varsha PV, Satyam S. Review on *Thymus vulgaris* traditional uses and pharmacological properties. Med Aromat Plants. 2014;3(4):167. doi: 10.4172/2167-0412.1000167.
- Liu W, Zhu Y, Wang Y, Qi S, Wang Y, Ma C, et al. Antiamnesic effect of extract and alkaloid fraction from aerial parts of *Peganum harmala* on scopolamine-induced memory deficits in mice. J Ethnopharmacol. 2017;204:95-106. doi: 10.1016/j.jep.2017.04.019.
- Gülçın İ, Oktay M, Kıreçcı E, Küfrevioğlu Öİ. Screening of antioxidant and antimicrobial activities of anise (*Pimpinella* anisum L.) seed extracts. Food Chem. 2003;83(3):371-82. doi: 10.1016/s0308-8146(03)00098-0.
- 64. Mushtaq A, Anwar R, Ahmad M. Memory enhancing effect of anise (*Pimpinella anisum*) with respect to its antioxidant activity in albino mice. J Anim Plant Sci. 2019;29(2):602-10.
- 65. Boskabady MH, Shafei MN, Saberi Z, Amini S. Pharmacological effects of *Rosa damascena*. Iran J Basic Med Sci. 2011;14(4):295-307.
- Kizil S, Haşimi N, Tolan V, Kilinc E, Karataş H. Chemical composition, antimicrobial and antioxidant activities of hyssop (*Hyssopus officinalis* L.) essential oil. Notulae Botanicae Horti Agrobotanici Cluj-Napoca. 2010;38(3):99-103. doi: 10.15835/nbha3834788.
- 67. Nahida, Ansari SH, Siddiqui AN. Pistacia lentiscus: a review

on phytochemistry and pharmacological properties. Int J Pharm Pharm Sci. 2012;4(4):16-20.

- Takooree H, Aumeeruddy MZ, Rengasamy KRR, Venugopala KN, Jeewon R, Zengin G, et al. A systematic review on black pepper (*Piper nigrum* L.): from folk uses to pharmacological applications. Crit Rev Food Sci Nutr. 2019;59(Suppl 1):S210-43. doi: 10.1080/10408398.2019.1565489.
- Usmani A, Khushtar M, Arif M, Siddiqui MA, Sing SP, Mujahid M. Pharmacognostic and phytopharmacology study of *Anacyclus pyrethrum*: an insight. J Appl Pharm Sci. 2016;6(3):144-50. doi: 10.7324/japs.2016.60325.
- Sujith K, Darwin CR, Sathish, Suba V. Memory-enhancing activity of *Anacyclus pyrethrum* in albino Wistar rats. Asian Pac J Trop Dis. 2012;2(4):307-11. doi: 10.1016/s2222-1808(12)60067-x.
- 71. Sanaie S, Nikanfar S, Yousefi Kalekhane Z, Azizi-Zeinalhajlou A, Sadigh-Eteghad S, Araj-Khodaei M, et al. Saffron as a promising therapy for diabetes and Alzheimer's disease: mechanistic insights. Metab Brain Dis. 2023;38(1):137-62. doi: 10.1007/s11011-022-01059-5.
- 72. Balakumbahan R, Rajamani K, Kumanan K. *Acorus calamus*: an overview. J Med Plants Res. 2010;4(25):2740-5.
- 73. Malik R, Kalra S, Pooja, Singh G, Meenu, Gahlot V, et al. Antioxidative and neuroprotective potential of *Acorus calamus* Linn. and *Cordia dichotoma* G. Forst. In Alzheimer's type dementia in rodent. Brain Res. 2024;1822:148616. doi: 10.1016/j.brainres.2023.148616.
- 74. Rasi Marzabadi L, Fazljou SM, Araj-Khodaei M, Sadigh-Eteghad S, Naseri A, Talebi M. Saffron reduces some inflammation and oxidative stress markers in donepeziltreated mild-to-moderate Alzheimer's disease patients: a randomized double-blind placebo-control trial. J Herb Med. 2022;34:100574. doi: 10.1016/j.hermed.2022.100574.
- Sharififar F, Dehghn-Nudeh G, Mirtajaldini M. Major flavonoids with antioxidant activity from *Teucrium polium* L. Food Chem. 2009;112(4):885-8. doi: 10.1016/j. foodchem.2008.06.064.
- Peterson DW, George RC, Scaramozzino F, LaPointe NE, Anderson RA, Graves DJ, et al. Cinnamon extract inhibits tau aggregation associated with Alzheimer's disease in vitro. J Alzheimers Dis. 2009;17(3):585-97. doi: 10.3233/jad-2009-1083.
- 77. Chnadrashekhar C, Sharma S, Mukopadayay S. A review on memory enhancing activity of ginger. Int J Health Sci. 2022;6(S1):14114-27. doi: 10.53730/ijhs.v6nS1.8576.
- Haghaei H, Soltani S, Arefhosseini SR, Rashidi MR, Karima S. Boswellic acids as promising leads in drug development against Alzheimer's disease. Pharm Sci. 2020;27(1):14-31. doi: 10.34172/ps.2020.25.
- 79. Ahmadian-Attari MM, Ahmadiani A, Kamalinejad M, Dargahi

L, Shirzad M, Mosaddegh M. Treatment of Alzheimer's disease in Iranian traditional medicine. Iran Red Crescent Med J. 2015;17(1):e18052. doi: 10.5812/ircmj.18052.

- 80. Srivastava P. Therapeutic potential of *Piper longum* L. for disease management-a review. Int J Pharma Sci. 2014;4(4):692-6.
- Sivapalan SR. Medicinal uses and pharmacological activities of *Cyperus rotundus* Linn-a review. International Journal of Scientific and Research Publications. 2013 May;3(5):1-8.
- Rabiei Z, Hojjati M, Rafieian-Kopaeia M, Alibabaei Z. Effect of *Cyperus rotundus* tubers ethanolic extract on learning and memory in animal model of Alzheimer. Biomed Aging Pathol. 2013;3(4):185-91. doi: 10.1016/j.biomag.2013.08.006.
- Gaire BP, Subedi L. Phytochemistry, pharmacology and medicinal properties of *Phyllanthus emblica* Linn. Chin J Integr Med. 2014. doi: 10.1007/s11655-014-1984-2.
- 84. Gupta A, Kumar R, Kumar S, Pandey AK. Pharmacological aspects of *Terminalia bellirica*. In: Mahdi AA, Abid M, Khan MM, Ansari MI, Maheshwari RK, eds. Molecular Biology and Pharmacognosy of Beneficial Plants. Delhi, India: Lenin Media Private Limited; 2017. p. 52-64.
- 85. Chaudhari GM, Mahajan RT. Comparative antioxidant activity of twenty traditional Indian medicinal plants and its correlation with total flavonoid and phenolic content. Int J Pharm Sci Rev Res. 2015;30(1):105-11.
- Parihar S, Sharma D. A brief overview on *Vitis vinifera*. Sch Acad J Pharm. 2021;12(12):231-9. doi: 10.36347/sajp.2021. v10i12.005.
- Ibrahim Fouad G, Zaki Rizk M. Possible neuromodulating role of different grape (*Vitis vinifera* L.) derived polyphenols against Alzheimer's dementia: treatment and mechanisms. Bull Natl Res Cent. 2019;43(1):108. doi: 10.1186/s42269-019-0149-z.
- Taheri-Targhi S, Gjedde A, Araj-Khodaei M, Rikhtegar R, Parsian Z, Zarrintan S, et al. Avicenna (980-1037 CE) and his early description and classification of dementia. J Alzheimers Dis. 2019;71(4):1093-8. doi: 10.3233/jad-190345.
- 89. Moini Jazani A, Farajpour Maleki R, Kazemi AH, Ghasemi Matankolaei L, Taheri Targhi S, Kordi S, et al. Intestinal helminths from the viewpoint of traditional Persian medicine versus modern medicine. Afr J Tradit Complement Altern Med. 2018;15(2):58-67. doi: 10.21010/ajtcamv15i2.8.
- Tajadini H, Choopani R, Saifadini R. Lifestyle methods for prevention of Alzheimer's disease from the perspective of traditional Persian medicine. J Evid Based Complementary Altern Med. 2016;21(3):243-5. doi: 10.1177/2156587215611787.
- Namiranian P, Karimi M, Emami Razavi SZ, Fateh Garoos A, Ayati MH. Visceral manipulation in resources of Persian medicine. Tradit Integr Med. 2021;6(2):158-65. doi: 10.18502/tim.v6i2.6795.